

Spectral curves for Cauchy–Riemann operators on punctured elliptic curves

C. Bohle ^{*} I.A. Taimanov [†]

1. Introduction. The spectral curve of a periodic differential operator is fundamental to finite gap integration theory. The solution of the inverse problem for a finite gap operator consists in assigning such an operator to the spectral data consisting of the spectral curve and some additional data related to it. For one–dimensional Schrödinger operators the spectral curve was defined in [1], and in [2] this notion was generalized to two–dimensional Schrödinger operators for which the spectral curve parametrizes the Floquet–Bloch functions at a fixed energy level.

In the present article we show that an analogous spectral curve can be defined for the Cauchy–Riemann operator on a punctured elliptic curve, provided one imposes appropriate boundary conditions.

Theorem 1 *Let X be an elliptic curve $X = \mathbb{C}/\{\mathbb{Z}e_1 + \mathbb{Z}e_2\}$, where e_1 and e_2 are generators of the period lattice, and let p_1, \dots, p_N be pair–wise distinct points on X . We consider the linear problem*

$$\bar{\partial}\psi = 0, \tag{1}$$

$$\psi(z + e_j) = e^{2\pi i \langle k, e_j \rangle} \psi(z), \quad j = 1, 2, \tag{2}$$

where $k = (k_1, k_2) \in \mathbb{C}^2$, $\langle k, e_j \rangle = k_1 \operatorname{Re} e_j + k_2 \operatorname{Im} e_j$ and ψ satisfies

$$\psi(z) = \frac{a_j}{z - p_j} + O(|z - p_j|) \text{ for } j = 1, \dots, N \tag{3}$$

for some $a_j \in \mathbb{C}$. The solutions to (1)–(3) with k and a_1, \dots, a_N arbitrary are parametrized (up to scale) by a one–dimensional (spectral) curve, see (7), which up to one puncture is an N –sheeted covering of X .

^{*}Universität Tübingen, Fachbereich Mathematik, Auf der Morgenstelle 10, 72076 Tübingen, Germany; e-mail:bohle@mathematik.uni-tuebingen.de

[†]Sobolev Institute of Mathematics, 630090 Novosibirsk, Russia; e-mail:taimanov@math.nsc.ru.

Algebraic curves of the type thus obtained appear as irreducible components of spectral curves of minimal tori with planar ends in \mathbb{R}^3 (see §4 below). It appears that these curves coincide with the spectral curves of certain elliptic KP solitons as studied by Krichever [3].

It should be noted that, unlike in the theory of parabolic bundles, we do not prescribe principal parts (as the a_j are not fixed), but cut down dimensions by demanding that the order zero terms in the asymptotic expansions (3) at the punctures vanish.

2. Spectral curves of Schrödinger operators. Let $L = \partial\bar{\partial} + U$ be a two-dimensional Schrödinger operator with a double-periodic potential: $U(z + e_1) = U(z + e_2) = U(z)$. We say that ψ is a Floquet eigenfunction of L with the eigenvalue E if it is a formal solution to the equation

$$L\psi = E\psi$$

which satisfies the periodicity conditions (2). The quantities k_1 and k_2 are called the quasimomenta of ψ and $(\nu_1, \nu_2) = (e^{2\pi i\langle k, e_1 \rangle}, e^{2\pi i\langle k, e_2 \rangle})$ are the Floquet multipliers of ψ .

The spectral curve which parametrizes the Floquet functions for $E = 0$ up to scale was defined in [2], where it was shown how to reconstruct an operator L from a (complex) curve of finite genus and some additional data. Such operators are called finite gap at the zero energy level.

In general spectral curves are not finite genus and there are two ways to rigorously establish their existence: by using the Fredholm alternative for analytic pencils of operators [4], or by perturbing the spectral curve for $L_0 = \partial\bar{\partial}$ [5].

3. Spectral curves of tori in \mathbb{R}^3 . The Weierstrass representation assigns to every solution $\psi: X \rightarrow \mathbb{C}^2$ of the problem

$$\mathcal{D}\psi = 0,$$

where \mathcal{D} is a Dirac operator with a double-periodic real-valued potential

$$\mathcal{D} = \begin{pmatrix} 0 & \partial \\ -\bar{\partial} & 0 \end{pmatrix} + \begin{pmatrix} U & 0 \\ 0 & U \end{pmatrix}, \quad U = \bar{U},$$

a conformal immersion of X or its covering into \mathbb{R}^3 , determined by

$$x^k = x^k(0) + \int \left(x_z^k dz + \bar{x}_z^k d\bar{z} \right), \quad k = 1, 2, 3,$$

with

$$x_z^1 = \frac{i}{2}(\bar{\psi}_2^2 + \psi_1^2), \quad x_z^2 = \frac{1}{2}(\bar{\psi}_2^2 - \psi_1^2), \quad x_z^3 = \psi_1\bar{\psi}_2, \quad (4)$$

branched at zeros of ψ and with a double-periodic Gauss map (see [6, 7]). Moreover, every conformal immersion of X does admit such a representation if one allows for ψ to have $\mathbb{Z}_2 = \{\pm 1\}$ -monodromy, i.e., to be the section of a spinor bundle on X [7].

Such a representation exists for all immersed surfaces [7], and for $U = 0$ it reduces to the classical Weierstrass representation of minimal surfaces.

The spectral curve of a torus in \mathbb{R}^3 is defined, in the spirit of [2], as the spectral curve of the corresponding Dirac operator \mathcal{D} at the zero energy level [4]. It parametrizes Floquet functions, taken up to scale, that satisfy the equation $\mathcal{D}\psi = 0$ and the conditions (2).

This spectral curve encodes the Willmore functional $\mathcal{W} = \int H^2 d\mu = 4 \int U^2 dx \wedge dy$, where H is the mean curvature of a closed oriented surface immersed into \mathbb{R}^3 and $d\mu$ is the measure induced by the immersion. Willmore tori, i.e. the critical points of the Willmore functional, are described by integrable systems. They split into several classes one of which are the stereographic projections of minimal tori with planar ends [8].

For Willmore tori that are not Euclidean minimal with planar ends it was proven in [9] that the spectral curve can be compactified by adding two points at infinity. (The result of [9] more generally concerns constrained Willmore tori, i.e. the critical points of \mathcal{W} restricted to tori within a fixed conformal class.)

4. Spectral curves of minimal tori with planar ends. The spectral curves of Willmore tori coming from minimal tori with planar ends have recently been described in [10] as reducible algebraic curves with two irreducible components Γ_+ and Γ_- which are complex conjugate to each other and exactly of the type arising in Theorem 1.

In terms of the Weierstrass representation minimal surfaces correspond to the trivial potential $U = 0$. Hence their spinor ψ splits into two components ψ_1, ψ_2 such that ψ_1 and $\bar{\psi}_2$ satisfy

$$\bar{\partial}\psi = 0.$$

By a planar end we mean a minimal conformal immersion $f : D \setminus \{p\} \rightarrow \mathbb{R}^3$ of a punctured disc $D \setminus \{p\}$ such that $\lim_{q \rightarrow p} f(q) = \infty$ and the immersion extends to an immersed disc in the conformal 3-sphere $S^3 = \mathbb{R} \cup \{\infty\}$. Such end p is characterized by $x_z^j, j = 1, 2, 3$ having poles of order 2 with vanishing residues. In terms of $\psi = (\psi_1, \psi_2)$ this condition becomes (3).

5. Proof of Theorem 1. It appears that Theorem 1 may be proven by a method very similar to the one used by Krichever [3].

To explicitly construct meromorphic functions satisfying (2), we use the basic elliptic functions $\sigma(z)$ and $\zeta(z)$ on $X = \mathbb{C}/\Lambda = \mathbb{C}/\{\mathbb{Z}e_1 + \mathbb{Z}e_2\}$ defined

by

$$\sigma(z) = z \prod_{m,n \in \mathbb{Z}^2 \setminus \{0\}} \left(1 - \frac{z}{e_{mn}}\right) \exp\left(\frac{z}{e_{mn}} + \frac{1}{2} \frac{z^2}{e_{mn}^2}\right), \quad e_{mn} = me_1 + ne_2,$$

$$\zeta(z) = \frac{d}{dz} \log \sigma(z).$$

It is known that

$$\sigma(z + e_j) = -\sigma(z)e^{\eta_j(z+e_j/2)}, \quad \zeta(z + e_j) = \zeta(z) + \eta_j, \quad j = 1, 2,$$

where $\eta_j = 2\zeta(e_j/2)$ satisfy $\eta_1 e_2 - \eta_2 e_1 = 2\pi i$. Following [3], we define

$$\Phi(z, \alpha) = \frac{\sigma(\alpha - z)}{\sigma(\alpha)\sigma(z)} e^{\zeta(\alpha)z}, \quad \alpha \in \mathbb{C} \setminus \Lambda$$

which is Λ -periodic in α , so that it's well defined for $\alpha \in X \setminus \{0\}$ (this is in fact the Baker–Akhiezer function for the one-gap Lamé potential).

For given multipliers $\nu_1 = e^{2\pi i \langle k, e_1 \rangle}$, $\nu_2 = e^{2\pi i \langle k, e_2 \rangle}$ (see (2)) which are not of the form $e^{\beta e_1}$, $e^{\beta e_2}$ there are unique $\alpha \in X \setminus \{0\}$ and $\mu \in \mathbb{C}$ such that any meromorphic function ψ with simple poles at $z_0 + \Lambda$ for which the periodicity conditions (2) hold coincides (up to scale) with

$$\Psi_{\mu, \alpha}(z - z_0) = e^{\mu z} \Phi(z - z_0, \alpha).$$

Riemann–Roch implies that, for given ν_1 , ν_2 and corresponding α , μ , any solution to the linear problem (1)–(2) with first order poles at the p_1, \dots, p_n has to be of the form

$$\psi = \sum_{l=1}^N a_l \Psi_{\mu, \alpha}(z - p_l). \quad (5)$$

The conditions (3) are then fulfilled if and only if

$$\mu a_l + \sum_{m \neq l} a_m \Phi(p_l - p_m, \alpha) = 0, \quad l = 1, \dots, N.$$

This linear system has a nontrivial solution $a = (a_1, \dots, a_N) \in \mathbb{C}^N$ if and only if its determinant vanishes. This condition

$$\det A(\mu, \alpha) = 0, \quad A_{lm} = \delta_{lm} \mu + (1 - \delta_{lm}) \Phi(p_m - p_l, \alpha) \quad (6)$$

yields an equation of the form

$$\mu^N + q_1(\alpha)\mu^{N-1} + \dots + q_N(\alpha) = 0. \quad (7)$$

The spectral curve Γ is given by the solutions to this equation. It is an N -sheeted covering of $X \setminus \{0\}$ with the natural projection $(\mu, \alpha) \in \Gamma \rightarrow \alpha \in X \setminus \{0\}$. The solutions ψ to (1)–(3) parametrized by Γ have Floquet multipliers

$$\nu_1 = \exp((\mu + \zeta(\alpha))e_1 - \alpha\eta_1) \quad \text{and} \quad \nu_2 = \exp((\mu + \zeta(\alpha))e_2 - \alpha\eta_2). \quad (8)$$

Note that precisely the same condition (6) (although in a different context) appears in [3], see (22) and the first half of (16) there assuming that $\dot{x}^k = 0$ for all k . In (25) of [3] it is shown that near $\alpha = 0$ the curve Γ has one sheet at which $\mu + \zeta(\alpha)$ has a pole at $\alpha = 0$ and $N - 1$ sheets at which the Floquet multipliers (8) converge to multipliers of the form $(\nu_1, \nu_2) = (e^{\beta e_1}, e^{\beta e_2})$ with β equal to the corresponding limit of $\mu + \zeta(\alpha)$ when α goes to 0.

Floquet functions ψ for multipliers of the form $(\nu_1, \nu_2) = (e^{\beta e_1}, e^{\beta e_2})$ cannot be obtained from the ansatz (5), but from

$$\psi = e^{\beta z} \left(a_0 + \sum_{l=1}^N a_l \zeta(z - p_l) \right) \quad (9)$$

with

$$\sum_{l=1}^N a_l = 0. \quad (10)$$

Condition (3) for this ansatz takes the form

$$a_0 + \beta a_k + \sum_{l \neq k} a_l \zeta(p_k - p_l) = 0, \quad k = 1, \dots, N.$$

After elimination of a_0 , these conditions together with (10) yield N linear equations for a_1, \dots, a_N . Taking the determinant yields a polynomial equation of order $N - 1$ for β . Its solutions correspond to the $N - 1$ ends of Γ for which the Floquet multipliers have a limit when α goes to 0.

EXAMPLE. In the simplest case $N = 1$ we have $\Gamma = X \setminus \{0\}$, the spectral curve of the one-gap Lamé potential.

ACKNOWLEDGEMENT. The first author (C.B.) was supported by DFG Sfb/Tr 71 ‘‘Geometric Partial Differential Equations’’, the second author

(I.A.T.) was supported by the program of the Presidium of RAS “Fundamental Problems of Nonlinear Dynamics in Mathematical and Physical Sciences”, in addition both authors were supported by Hausdorff Institute of Mathematics in Bonn.

References

- [1] Novikov, S.P.: A periodic problem for the Korteweg-de Vries equation. I. *Functional Anal. Appl.* **8**:3 (1974), 236–246.
- [2] Dubrovin, B.A., Krichever, I.M., and Novikov, S.P.: The Schrödinger equation in a periodic field and Riemann surfaces. *Soviet Math. Dokl.* **17** (1976), 947–952.
- [3] Krichever, I.M.: Elliptic solutions of the Kadomtsev-Petviashvili equation and integrable systems of particles. *Funct. Anal. Appl.* **14**:4 (1980), 282–290.
- [4] Taimanov, I.A.: The Weierstrass representation of closed surfaces in \mathbb{R}^3 . *Funct. Anal. Appl.* **32**:4 (1998), 258–267.
- [5] Krichever, I.M.: Spectral theory of two-dimensional periodic operators and its applications, *Russian Math. Surveys* **44**:2 (1989), 145–225.
- [6] Konopelchenko, B.G.: Induced surfaces and their integrable dynamics. *Stud. Appl. Math.* **96** (1996), 9–52.
- [7] Taimanov, I.A.: Modified Novikov–Veselov equation and differential geometry of surfaces. *Amer. Math. Soc. Transl., Ser. 2, V.* 179, 1997, pp. 133–151.
- [8] Bryant, R.L.: A duality theorem for Willmore surfaces. *J. Differential Geom.* **20** (1984), 23–53.
- [9] Bohle, C.: Constrained Willmore tori in the 4-sphere. *J. Differential Geom.* **86** (2010), 71–131.
- [10] Bohle, C., and Taimanov, I.A.: Euclidean minimal tori with planar ends and elliptic solitons, in preparation.